

Spectral universality of strong shocks accelerating charged particles

M.A. Malkov

Max-Planck Institut für Kernphysik, D-69029, Heidelberg, Germany

As a rule, the shock compression controls the spectrum of diffusively accelerated particles. We argue that this is not so if the backreaction of these particles on the shock structure is significant. We present a self-similar solution in which the accelerated particles change the flow structure near the shock so strongly that the total shock compression may become arbitrarily large. Despite this, the energy spectrum behind the shock is close to $E^{-3/2}$ independently of anything at all.

52.35.Tc, 95.30.Lz, 98.70.Sa

The first order Fermi acceleration at shocks was always regarded as one of those universal phenomena that result in a scale-invariant behavior of physical quantities. In particular, this acceleration mechanism has predicted a power-law momentum distribution [1,2] similar to those observed in diverse astrophysical objects. There exists, however, another important characteristics, the energy content of accelerated particles. Generally speaking, this quantity makes the expectations of the scale invariance not so well grounded. Whenever the shock accelerates efficiently in the sense of its energy conversion into energetic particles (and this is precisely what one anticipates in the context of the problem of cosmic ray origin), a more complicated flow structure must form in place of this shock. Besides the shock itself, usually called subshock, it contains a long precursor in which the inflowing gas is gradually decelerated by the pressure gradient of energetic particles, in other words of the cosmic rays (CRs) produced in the entire shock transition, Fig. 1. One interesting aspect of this gas-CR coupling is an increase of the total compression ratio which should harden the spectrum and make the CR acceleration potentially much more efficient. At the same time, the most attractive feature of this acceleration mechanism, the scale-free form of the spectrum is likely to be lost since the shock develops now its internal structure and particles with different momenta “see” different shock compression due to the strong momentum dependence of the CR diffusivity. It is also well known (see e.g., Ref. [2]) that this broadening of the shock transition softens the spectrum. The combined effect of these two oppositely acting factors has been resisting any regular analytic calculations for a long time (see, however, [3] for a perturbative approach).

Recently [4], we have reduced the problem of the gas-CR coupling to one nonlinear integral equation. Its solution does provide selfconsistently both the particle spectrum and the structure of the hydrodynamic flow. At least in a certain phase space region this solution turns

out to preserve the scale-invariance of the test particle results. But the most surprising aspect of this solution is that in contrast to the well known linear solution for the particle momentum distribution $f \propto p^{-q}$ where $q = 3r/(r-1)$ and r stands for the shock compression ratio, in the nonlinear regime $q = 3\frac{1}{2}$ independently of anything at all. This means that when accelerated particles modify the shock strongly their spectrum and the flow profile mutually adjust in such a way that *all such shocks become universal accelerators producing essentially the same power-law spectra*. They differ only in the cut-off momentum that is always limited by the size and/or the age of the shock. The above result is, however, limited formally to relativistic particles with the Bohm diffusivity $\kappa \propto p$ and relates to the solution (out of the three possible) with the highest conversion efficiency of the flow energy to the CR energy.




FIG. 1. The flow structure in a strongly modified CR shock.

In this *letter*, using a simpler mathematical technique, we demonstrate that the spectrum universality persists also in the nonrelativistic region and for even more general form of diffusion coefficient, e.g., $\kappa \propto p^2 / \sqrt{1 + p^2/m^2 c^2}$. The power-law index, being $q = 3\frac{1}{2}$ in the region $p \gg mc$, simply changes for $q = 4$ in the region $p \ll mc$ before it conforms to the local subshock compression r_s , i.e. approaches $q = 3r_s/(r_s - 1)$ at $p \sim p_0 \ll mc$, where p_0 is the injection momentum [4,5]. Remarkably, the $q = 3\frac{1}{2}$ index coincides with the plain test particle result for a strong shock in a purely thermal relativistic gas, whereas $q = 4$ coincides with that for a nonrelativistic gas. All these in spite of the facts that the flow profile is strongly modified and the total compression ratio may be much higher than 4 ($q = 4$) and even 7 ($q = 3\frac{1}{2}$) occurring in nonrelativistic and relativistic gases, respectively. Moreover, the subshock compression

ratio r_s may be significantly lower than 4 and cannot be responsible for $q = 4$ spectrum in the nonrelativistic momentum range.

The standard formulation of the problem includes the diffusion-convection equation constrained by conservation of the fluxes of mass and momentum (see e.g., Ref. [2]). We assume that a strong CR-modified shock propagates in the positive x -direction in a cold medium with the mass density ρ_1 . In the shock frame of reference the steady mass flow profile is defined as follows $U(x) = -u(x)$, $x \geq 0$ and $U(x) = -u_2$, $x < 0$, where u_2 is the (constant) downstream mass velocity, $u(0+) = u_0$, and $u(x) \rightarrow u_1 = \text{const}$, as $x \rightarrow \infty$, Fig. 1. The equations read

$$\frac{\partial}{\partial x} \left(u g + \kappa(p) \frac{\partial g}{\partial x} \right) = \frac{1}{3} \frac{du}{dx} p \frac{\partial g}{\partial p}, \quad (1)$$

$$\rho u = \rho_1 u_1, \quad (2)$$

$$P_c + \rho u^2 = \rho_1 u_1^2, \quad x > 0 \quad (3)$$

Here the number density of CRs is normalized to $4\pi g dp/p$. The upper limit p_1 stands for a boundary in the momentum space (cut-off) beyond which particles are assumed to leave the system instantaneously ($g \equiv 0$, $p > p_1$). In the downstream medium, $x < 0$, the only bounded solution is $g = G(p) \equiv g(p, x = 0)$. In Eqs. (2,3) $\rho(x)$ is the mass density, $\rho_1 = \rho(\infty)$, P_c is the CR pressure

$$P_c(x) = \frac{4\pi}{3} mc^2 \int_{p_0}^{p_1} \frac{p dp}{\sqrt{p^2 + 1}} g(p, x) \quad (4)$$

The particle momentum p is normalized to mc . As indicated, Eq. (3) is written in the region $x > 0$ where we have neglected the contribution of the adiabatically compressed cold gas (i.e., particles with $0 < p < p_0$) confining our consideration to sufficiently strong shocks with $M^2 \equiv \rho_1 u_1^2 / \gamma P_{g1} \gg (u_1/u_0)^\gamma$, where γ is the specific heat ratio of the plasma and P_{g1} is the gas kinetic pressure far upstream, i.e., at $x = \infty$ (see [4] for a detailed discussion of this approximation). The subshock strength can be obtained from the familiar Rankine-Hugoniot condition for the gas

$$r_s \equiv \frac{u_0}{u_2} = \frac{\gamma + 1}{\gamma - 1 + 2(u_1/u_0)^{\gamma+1} M^{-2}} \quad (5)$$

Introducing the flow potential ϕ , such as $u = d\phi/dx$ we seek the solution of Eq. (1) in the form

$$g = g_0(p) \exp \left\{ -\frac{1+\beta}{\kappa(p)} \phi \right\}, \quad x > 0 \quad (6)$$

where

$$\beta(p) \equiv -(1/3) d \ln g_0 / d \ln p \quad (7)$$

This is the key substitution in our analysis. Considering u as $u(\phi)$, substituting Eq.(6) in Eq. (1) and separating the variables we obtain

$$\phi \frac{du}{d\phi} = \lambda u \quad (8)$$

$$p \frac{d\beta}{dp} = (1 + \beta) \left(\frac{d \ln \kappa}{d \ln p} - \frac{3}{\lambda} \beta \right) \quad (9)$$

where λ is a constant. Eq. (8) may be readily integrated and yields for the flow potential

$$\phi(x) = \phi_0^{-\lambda/(1-\lambda)} [(1-\lambda)u_0 x + \phi_0]^{1/(1-\lambda)} \quad (10)$$

where $\phi_0 = \phi(0)$ is another constant (see below).

The solution for the particle spectrum $g_0(p)$ to be obtained from the system (7,9) depends on two further integration constants. These are the magnitude and the slope of g_0 at the injection momentum $p = p_0$, i.e., $g_0(p_0)$ and $\beta_0 \equiv \beta(p_0)$, respectively, provided by an “injection” solution [5]. Throughout this paper we shall consider $g_0(p_0)$ as given. The constant β_0 may also be determined from merely the subshock strength (see also ref. [4] and below). It is straightforward to verify that the following expression is the first integral of the system (7,9)

$$g_0(p) \left(\frac{\kappa}{1+\beta} \right)^\lambda = g_0(p_0) \left(\frac{\kappa_0}{1+\beta_0} \right)^\lambda \quad (11)$$

where $\kappa_0 \equiv \kappa(p_0)$. The solution for g_0 is given by

$$g_0(p) = g_0(p_0) \left(\frac{p}{p_0} \right)^3 \left[1 + 3 \frac{\beta_0 + 1}{\lambda \kappa_0} p_0^{-3/\lambda} \int_{p_0}^p \kappa(p') p'^{3/\lambda-1} dp' \right]^{-\lambda} \quad (12)$$

For p not too close to p_0 , more precisely for $(\kappa/\kappa_0)(p/p_0)^{3/\lambda} \gg 1$, the particle spectrum g_0 acquires a form determined by merely that of $\kappa(p)$ and “forgets” its behavior at $p \sim p_0$. Note that the latter (and thus the parameter β_0) is prescribed by the subshock strength, according to the relation $d \ln G / d \ln p = 3/(r_s - 1)$ at $p = p_0$, which can be recast with the help of (11) as

$$\beta_0 - \phi_0 \beta_0 \frac{1 + \beta_0}{\kappa_0} = \frac{1}{r_s - 1} \quad (13)$$

If p is large in the above sense and $\kappa(p)$ is a power-law, then the spectrum $g_0(p)$ must also be, $g_0(p) \propto \kappa^{-\lambda}(p)$ and

$$\beta \simeq (\lambda/3) d \ln \kappa / d \ln p \quad (14)$$

(the last formula may be also obtained from Eq.(9) assuming $\beta \simeq \text{const}$). As we shall see, the parameter λ depends on the scaling of κ as well, and the most surprising consequence of this dependence is that the resulting slope of $g_0(p)$ is, in fact, independent of $\kappa(p)$.

What we obtained so far is a one parameter family of exact solutions to equation (1) that require a rather special class of the flow profiles $u(\phi)$. It is by no means guaranteed that any of these solutions satisfy the Bernoulli's

integral (3) (the continuity condition (2) can be easily satisfied since it does not depend on $g(p > p_0)$). Indeed, the only unspecified quantity in solution (6,10,12) that can potentially be used to satisfy the functional relation (3), is the constant λ which is, generally speaking, not enough. Fortunately, the dependence of $P_c(\phi)$ in Eq.(3) can be made consistent with the dependence $u(\phi) \propto \phi^\lambda$, following from Eq.(8), in a fairly large part of shock transition, provided that the parameter λ is chosen properly. To demonstrate this we substitute Eq.(6) into Eqs.(3,4). Using Eq.(2), the Bernoulli's integral (3) rewrites

$$u(\phi) + \zeta \int_{s_1}^{s_0} \frac{ds}{\beta(s)} s^{\lambda-1} e^{-\phi s} \frac{p^2(s)}{\sqrt{1+p^2(s)}} = u_1 \quad (15)$$

where we have introduced a new variable s in place of p

$$s = \frac{1+\beta}{\kappa} \quad (16)$$

and the limits $s_{0,1} = s(p_{0,1})$. We have also used the first integral (11), $g_0 \propto s^\lambda$. The injection rate ζ is defined here as

$$\zeta = \frac{4\pi\lambda}{9} \frac{mc^2}{\rho_1 u_1} \left(\frac{\kappa_0}{1+\beta_0} \right)^\lambda g_0(p_0) \quad (17)$$

and the function $p(s)$ in Eq.(15) should be determined from Eq.(16). We now make a “standard” assumption: $\kappa(p) = Kp^2(1+p^2)^{-1/2}$, i.e., the mean free path of a particle is proportional to its Larmor radius (here K is a reference diffusivity). Then, Eq.(15) rewrites

$$u(\phi) = u_1 - \frac{\zeta}{K} \int_{s_1}^{s_0} \left(1 + \frac{1}{\beta(s)} \right) s^{\lambda-2} e^{-\phi s} ds \quad (18)$$

According to Eqs.(14,16), $\beta(s)$ is a very simple function, taking in the most part of its domain nearly constant and relatively close values, $\beta \simeq \lambda/3$ for $Ks \ll 1$ and $\beta \simeq 2\lambda/3$ for $Ks \gg 1$. It changes monotonically between these limiting values. Therefore, the last integral allows a straightforward asymptotic analysis. Our goal here is to examine whether Eq.(8), according to which $du/d\phi \propto \phi^{\lambda-1}$ is compatible with Eq.(18). We assume that $0 < \lambda < 1$, which will be confirmed later. Differentiating Eq.(18) with respect to ϕ and considering first the region $1/s_0 \ll \phi \ll 1/s_1$ we may obviously replace the lower limit by zero and the upper one by infinity. From Eq.(18) we then obtain

$$\begin{aligned} \frac{du}{d\phi} &\simeq \frac{\zeta}{K\phi^\lambda} \int_0^\infty [1 + 1/\beta(\tau/\phi)] \tau^{\lambda-1} e^{-\tau} d\tau \\ &= \frac{\zeta\Gamma(\lambda)}{K\phi^\lambda} [1 + 1/\beta(\bar{\tau}/\phi)] \end{aligned} \quad (19)$$

where Γ is the gamma function and $\beta(\tau/\phi)$ is replaced in the last integral by its mean value at $\bar{\tau}/\phi$ with $\bar{\tau} \sim 1$. As we have seen already the function $\beta(\bar{\tau}/\phi)$ in the last

expression varies slowly and it is close to $\lambda/3$ for $\phi > K$ and to $2\lambda/3$ for $\phi < K$. Therefore, the ϕ dependence of $du/d\phi$ is determined by the factor $\phi^{-\lambda}$ and is indeed consistent with Eq.(8), i.e., with $u \propto \phi^\lambda$ for $\lambda = 1/2$. Eq.(19) becomes invalid for $\phi \gtrsim 1/s_1$, since the lower limit in Eq.(18) cannot be replaced by zero in this case. The function $du/d\phi$ cuts off as (see Eq.(18))

$$\frac{du}{d\phi} \simeq \frac{\zeta}{K\sqrt{s_1}\phi} \left(1 + \frac{1}{\beta(s_1)} \right) e^{-s_1\phi} \quad (20)$$

The last asymptotics obviously matches the formula (19) at $\phi \sim 1/s_1$ but it is no longer consistent with Eq.(8). At the same time, this is a periphery of the shock transition where $u(\phi)$ exponentially approaches its limit u_1 at $x \sim l \equiv \kappa(p_1)/u_1$, Fig. 1. This particular form of the cut-off of $du/d\phi$ is entirely due to our assumption about an abrupt cut-off of the particle spectrum at $p = p_1$ ($s = s_1$ in Eq.(18)). Therefore, there is no major physical reason to refine our solution in this region unless a more realistic model of particle losses in the region $p \sim p_1$ is adopted. According to Eq.(10) the flow profile in the internal part of the precursor is simply linear ($\lambda = 1/2$)

$$u(x) = u_0 + \frac{u_0^2}{2\phi_0} x \quad (21)$$

An asymptotic approach resulting in a flow profile $u(x)$ that is uniformly valid (for all $x > 0$) and coincides with eq.(21) for $0 < x < l$ has been developed in Ref. [4]. It produces a full set of Rankine-Hugoniot relations, determining the total (u_1/u_2) and the subshock (u_0/u_2) compression ratios along with the constant ϕ_0 in Eq.(10) as functions of the Mach number, the injection rate and the upper cut-off momentum p_1 . These quantities have been calculated in Ref. [4] with a help of nonlinear integral equation derived from the system (1-3). This equation has three different solutions in a certain parameter range (see also [6]) with dramatically different acceleration efficiencies [7]. However, its derivation utilizes a relativistic form $\kappa \propto p$ for all p and a similar form of the pressure integral. These assumptions oversimplify the particle spectrum at $p < 1$ and we have dropped them here. According to the form of the particle spectrum with the index $q \equiv 3(\beta+1) = 3.5$ at high momenta $p > 1$ and the linear profile of the flow velocity (21) the solution obtained in the present paper can be identified with that of the Ref. [4] having the highest efficiency of CR production. Note that this solution becomes unique for sufficiently large ζ and p_1 . In addition, it provides parameters that have not been determined in the present treatment. First of all the relation between the flow deceleration in the precursor and in the subshock was found to be $(u_1 - u_0)/(u_0 - u_2) \propto \zeta p_1$ which, in combination with Eq.(5) determines the unknown velocities u_0 and u_2 given the far upstream velocity u_1 . Also the constant ϕ_0

may be related to the precursor length $l = \kappa(p_1)/u_1$ by $u_0^2/2\phi_0 \simeq u_1/l$.

As we already emphasized the scale-invariance does not exist everywhere in the phase space as in any other solution of this kind applied to a real physical situation. However, the scale invariant behavior in the region $u_0 < u < u_1$, $p_0 < p < p_1$ may span decades in both momentum and coordinate space in those astrophysical situations where the dynamical range p_1/p_0 is large. E.g., for typical supernova shock conditions one may expect $p_1/p_0 \sim 10^8$. It is also interesting to note that the downstream particle spectrum given by Eq.(11) with the approximate formula (14) for β and being expressed in terms of kinetic energy E rather than momentum exhibits a particularly uniform behaviour throughout the entire energy range, relativistic and nonrelativistic. In a standard normalization $F(E)dE$ this spectrum has the form

$$F \propto E^{-3/2} \sqrt{\frac{(E+1)(7E^2+14E+8)}{(E+2)^3}}$$

where E is measured in mc^2 , and the spectral index is close to 1.5 for all but injection energies (if the subshock is substantially reduced, see text below Eq.(12)). We emphasize that its value does not depend on any parameters involved in these calculations. This results from the mutual adjustment of the flow profile $u(x)$ and the particle distribution in energy and in this sense the form of the spectrum is completely universal.

One may ask then what does the power-law index depend on? It should be primarily the momentum dependence of the CR diffusivity $\kappa(p)$ that we have specified in our treatment above. To examine this idea we rescale κ as follows $\kappa' = \kappa^\alpha$, where $\alpha > 1/2$ and the rescaling of all primed variables below is induced by the above transformation of κ . According to Eq.(14) the spectral slope β is now to be replaced by $\beta' = (\alpha\lambda'/3)d \ln \kappa/d \ln p$. Recalculating $du/d\phi$ in Eq.(19) with these rescaled spectrum and CR diffusivity κ' we obtain $du/d\phi \propto \phi^{1/\alpha-1-\lambda'}$. Since the formula $u \propto \phi^{\lambda'}$ (Eq.(8)) holds, we deduce that the rescaled $\lambda' = 1/2\alpha$. Therefore the spectral slope β remains actually unchanged, $\beta' = \beta$.

We conclude that the spectral universality is insensitive to the scaling of the CR diffusivity $\kappa(p)$, in other words, to the spectrum of the underlying MHD turbulence. On the other hand, the velocity profile does depend on α . From the above analysis we obtain $u \propto x^{1/(2\alpha-1)}$ which also explains the condition $\alpha > 1/2$ [8]. There is a small deviation from this scaling at the distances x corresponding to the diffusion length of particles with $p \sim 1$ since β depends on ϕ in Eq.(19) at $\phi \sim K$.

Summarizing the last results, when $\kappa(p)$ rescales, so does the flow profile $u(x)$ but β remains invariant. It is not difficult to understand why this is so. As usual in the Fermi process the spectral slope of course depends on the

flow compression. But, since the flow is modified, a particle with momentum p , samples not the total compression but only a compression accessible to it. This compression is determined by the relation $\phi(x) \propto \kappa^\alpha$ (Eq. (6)). As we have shown, $u(\phi) \propto \phi^{1/2\alpha}$. Therefore, the flow compression, as seen by this particle diffusively bound to the shock front, scales as $u/u_2 \propto \phi^{1/2\alpha} \propto \sqrt{\kappa(p)}$. As this is independent of α the index β must also be.

This work was done within the Sonderforschungsbereich 328 of the Deutsche Forschungsgemeinschaft (DFG).

-
- [1] G.F. Krinsky, Dokl. Akad. Nauk SSSR **234**, 1306 (1977) (Engl. Transl. Sov. Phys.-Dokl. 23, 327); W. I. Axford, E. Leer, and G. Skadron, Proc. 15th ICRC **11**, 132 (Plovdiv, 1977); A. R. Bell, MNRAS **182**, 147; 443 (1978); R. D. Blandford and J. P. Ostriker, Astrophys. J. **221**, L29 (1978)
 - [2] Drury, L. O'C. 1983, Rep. Prog. Phys. 1983, 46, 973
 - [3] Blandford, R. D. 1980, Astrophys. J. , 238, 410
 - [4] M. A. Malkov, Astrophys. J. **485**, 638 (1997)
 - [5] M. A. Malkov and H. J. Völk, Astronomy and Astrophys. **300**, 605 (1995)
 - [6] M. A. Malkov, Astrophys. J. **491**, 584 (1997)
 - [7] The multiplicity of solutions for nonlinearly accelerating shocks is not new. It has been studied in the framework of a hydrodynamic description (see [2] and the references therein). However, there are very important differences between kinetic solutions obtained in [4], [6] and their hydrodynamic counterparts mostly because the latter are limited to the case in which the particle diffusivity κ is independent of momentum whereas the approaches in [4] and in the present paper not only allow for but also exploit this dependence.
 - [8] Remarkably, a precisely opposite condition $\alpha < 1/2$ is required to produce a steady velocity jump *without* momentum cut-off and injection but with a secularly broadening CR precursor. This has been shown by Drury [2]. The fact that our strictly stationary solution that is essentially based on energy losses through the upper cut-off and particle injection at $p = p_0$ appears immediately beyond $\alpha = 1/2$ is perhaps more than a coincidence. Also, the time saving numerical solutions with $\alpha \leq 1/2$ might be inadequate for modelling the more realistic case of $\alpha = 1$.